



GASTRO-INTESTINAL CLINICAL GENOMICS TEST MENU

GASTRO-INTESTINAL

| Sl No | TEST NAME | METHOD | SAMPLE TYPE | TAT |
|-------|---|--------|-----------------|---------|
| 1 | Wilson's disease gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 2 | Gilbert's/Crigler-Najjar syndrome gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 3 | Progressive familial intrahepatic cholestasis (PFIC) gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 4 | Primary Hyperoxaluria (Type I & II) | NGS | EDTA Blood 3 ml | 4 Weeks |
| 5 | Pancreatic Gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 6 | Glycogen storage disease gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 7 | Hemochromatosis Panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 8 | Alagille syndrome gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 9 | Polycystic liver disease gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 10 | Congenital Hepatic fibrosis gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |
| 11 | Tyrosinemia gene panel | NGS | EDTA Blood 3 ml | 4 Weeks |